

Dune Dynamics at DS '07

By Barry A. Cipra

The answer, my friend, is blowin' in the wind.

—Bob Dylan

Sand dunes are ubiquitous, sometimes problematic features of deserts and coastlines. Their undulations call to mind ripples on water, but dunes and water waves differ in crucial ways. In particular, whereas “traveling” water waves are largely illusory, a product of motion that is essentially up and down, the advance of sand dunes really does involve material transport: Individual grains of sand are blown by the wind along nearly horizontal trajectories, as anyone whose face has been sandblasted at the beach on a breezy day will know. Errant dunes have been known to block roads and overrun buildings.

Because of how they move, sand dunes also differ from water waves in shape. Of particular interest is a class of crescent-shaped dunes known as “barchans” (see Figure 1, top). In an invited address at this year’s SIAM Conference on Applications of Dynamical Systems, held in Snowbird, May 28 to June 1, Hans Herrmann of ETH Zürich described analyses he and colleagues have done of these fascinating formations.

The seminal scientific study of sand dunes, *The Physics of Blown Sand and Desert Dunes*, by Ralph Bagnold, dates back to 1941. In his book, Bagnold provided a Goldilocks-like classification for the aeolian (wind-driven) transport of sand. Sand that’s too coarse creeps along the ground; sand that’s too fine stays in suspension, blowing great distances. In the “just-right” phase, corresponding to grain diameters between 0.1 and 0.3 millimeters, sand moves by a process called “saltation,” from the Latin verb *saltare*, meaning “to leap.”

In saltation, the impact of a given sand grain transfers momentum to other grains,

which follow ballistic trajectories accelerated by the wind until they too hit the surface, kicking up other grains in turn. Because some kinetic energy is lost in the (inelastic) collisions, saltation dies out if the wind speed is too low, but it grows exponentially if the wind speed exceeds a certain threshold (see Figure 2). The latter case is self-limiting: Blown sand sucks energy out of the wind, eventually reducing its speed.

Saltation, Bagnold found, is the dominant mechanism in the formation of all sand dunes. The dunes will be barchans when wind blows mainly from one direction over largely flat terrain with relatively little sand. (When sand is abundant, formations known as transverse dunes are the rule.) Sand builds up on the backside of an incipient bump and slips around the sides to form a pair of arms reaching downwind. The build-up near the crest eventually reaches an unstable slope on the downwind side known as the angle of repose, which for sand is roughly 34 degrees. Sandslides leave the dune with an axial profile consisting of a shallow curve on the upwind side and a steep straight line downwind (see Figure 1, bottom).

Barchans typically appear in clusters, often made up of thousands of separate dunes of various sizes. Much like glaciers, barchans appear to be stationary but in fact advance slowly, on the order of centimeters per day. (Unlike glaciers, though, the physical transport of sand grains takes place only in a surface layer; a given grain travels in fits and starts, going great distances in a matter of days before being stochastically buried for years, decades, or centuries.) Small dunes, it’s been observed, travel faster than large ones. This begs the question, What happens when a small barchan overtakes a larger one?

People have not been measuring sand dune migration long enough to have observed the dynamics of interacting




Figure 2. Saltation dies out if wind speed is too low (solid curve), but becomes self-sustaining for winds in excess of a critical threshold (dashed curve). Figure courtesy of Hans Herrmann.

dunes—no ancient astronomers of sand kept meticulous records of barchan vagaries. Modern researchers have proposed various explanations, ranging from simple, permanent coalescence to soliton-like behavior in which the small dune merges momentarily with its bigger brother, only to re-emerge in its original size and shape on the other side.

Herrmann and colleagues, including current and former students Gerd Sauermann, Veit Schwämmle, and Orenicio Durán, have spent much of the current decade bringing computational precision to the study of dunes. They have extended the earlier analytic models of sand physics, developed computer models to solve the equations of shape and motion, and done field experiments with dunes in Morocco and Brazil, among other locations. Their *in silico* bar-

chans look a lot like the *in silico* originals.

An obvious starting point for analyzing sand flow is conservation of mass: The piling up or erosion of a dune at any point—i.e., the time derivative of dune height—is proportional to the gradient of the flux. The latter is a function of wind speed (itself a function of height) and the dynamics of saltation, which includes coefficients of drag and restitution. Because all motion occurs at the surface, the units of flux are kilograms per meter per second, rather than kilograms per meter squared per second; the constant of proportionality is the density of sand in the mobile surface layer.

Bagnold derived a simple formula for the flux, finding it to be proportional to the cube of the wind friction speed u , for speeds in excess of the threshold speed u_c , which is

See *Dunes* on page 16

SIAM News Story Helps Uncover a Copy of Green’s Essay of 1828

By Robert E. O’Malley, Jr.

We don’t know much about George Green (1793–1841). What we do know is found in Mary Cannell’s eloquent biography, published by SIAM shortly after her death in 2000. It was my pleasure to know this elegant Englishwoman, who devoted her retirement years to learning about Green and his impact.

Green was a miller in Sneinton, near Nottingham, who taught himself continental mathematics, presumably by reading works of Poisson, Laplace, and others from Bromley House, the Nottingham Subscription Library, which he joined in 1823. His famous work of 1828, “An Essay on the Application of Mathematical Analysis to the Theories of Electricity and Magnetism,” was printed by T. Wheelhouse of Nottingham for the author. According to Cajori [3], about 100 copies were printed. Fifty-one were sold at a price of 7 s., 6 d. (the weekly wage of a Nottingham stocking maker) to a list of subscribers obtained through advertisements in the *Nottingham Review* and *Nottingham Journal* in December 1827. In her biography of Green, Cannell [4] reports that local “clergymen, doctors, and the prosperous lace and textile manufacturers and merchants . . . made up more than half the sub-



scribers.” The book was also available from a few booksellers in Nottingham, Cambridge, and London.

The Essay is best known as the source of Green’s theorem (cf. Cross [6] and Archibald [1]), the electric potential, and the kernel now called a Green’s function (cf. Schwinger [8] and Challis and Sheard [5]). Green’s preface ends with the paragraph:

“Should the present Essay tend in any way to facilitate the application of analysis to one of the more interesting of the physical sciences, the author will deem himself amply repaid for any labour he may have bestowed upon it; and it is hoped the difficulty of the subject will incline mathematicians to read the work with indulgence, more particularly when they are informed that it was written by a young man, who has been obliged to obtain the little knowledge he possesses, at such intervals

See *Green’s Essay* on page 8




Figure 1. Anatomy of a dune. Forces exerted by wind, gravity, and friction determine the shape of a barchan. Figure courtesy of Hans Herrmann.

Dunes

continued from page 1

about a third of a meter per second. (Wind friction speed is derived from the vertical profile of wind speed, which depends on surface roughness.) In the 1970s, Heinz and Katharina Lettau of the University of Wisconsin modified Bagnold's formula to make flux proportional to $u^2(u - u_t)$. Herrmann's group has derived a yet more complicated formula based on a closer look at the details of saltation. Their formula still has flux asymptotic to u^3 for large u , but takes a different form for speeds just above the threshold and gives results that, after parameter fitting, agree well with wind tunnel experiments.

Using their equations and the commercial code FLUENT, Herrmann and colleagues have simulated barchan formation and motion. The computations generate the graceful crescent shapes of observed dunes and reproduce their relative migratory speeds. In a 2003 paper in *Nature*, Herrmann and Veit Schwämmle described soliton-like behavior when a smaller barchan catches up with a larger one. In further work, Herrmann and Durán found a range of behavior, depending on the ratio of dune volumes (dividing the smaller by the larger). Solitonic pass-through occurs when the ratio is larger than about 0.25. For ratios under 0.07, the smaller barchan is simply swallowed as it climbs the backside of the larger one. Between 0.07 and 0.14, the smaller dune emerges as multiple barchanettes, a process the researchers call "breeding." From 0.14 to 0.25, the smaller dune splits in two, as if "budding" off the horns of the larger barchan (see Figure 3).

The speed at which a barchan moves, Bagnold found, is inversely proportional to its height. That would seem to imply a nonsensically near-infinite speed for nanobarchans. In fact, barchans below a certain critical size are simply not seen. Herrmann, Durán, and Eric Parteli have shown that the barchan-forming threshold depends on the wind speed and the flux of sand in the area between dunes: As the values of these variables increase, the threshold drops. (Intuitively, this makes sense: A dune cannot move faster than the wind that carries the sand.)

Herrmann and Durán also recently derived a set of differential equations describing the competition between dune mobility and plant growth. Vegetation tends to retard erosion (mainly by slowing down the wind) and to persist until deposition buries (and kills) it. Plants take root most easily where the erosion/deposition rate is small to begin with. For barchans, this includes the horns of the crescent. As a consequence, the horns are pinned by plants while the crest between them advances, until the barchan inverts into an upwind-pointing parabola. Herrmann and Durán's equations offer insight into the saltation/vegetation battle. According to simulations based on their model, for example, inversion occurs only when the (dimensionless) ratio of characteristic erosion rate to vegetation growth velocity is smaller than about 0.5.

In another recent paper, Herrmann, Schwämmle, Durán, and Pedro Lind considered the dune size and spatial distributions in barchan fields. Their key equation, derived by a mean-field approach, relates




Figure 4. Alternating explanations. Unusual dunes on Mars (top) can be reproduced in simulations by assuming an alternating wind field (bottom). From "Saltation Transport on Mars," by Eric Parteli and Hans Herrmann. (Martian photos taken by the Mars Orbiter Camera, courtesy of NASA/JPL/MSSS.)

the characteristic interdune spacing L to the average dune width W and the standard deviation S of the distribution of widths: L is proportional to S^3/W^2 . In other words, the sparser the dune field, the greater the variability in dune size. (The size distribution itself is found to be log normal.)

Not unique to Earth, sand dunes are also abundant on Mars. Many of the crucial constants—gravity, wind resistance, and grain size—are different, but the physics is otherwise the same. In a detailed analysis, Herrmann and Parteli showed that saltation is an order of magnitude more sensitive to wind speed increments on Mars than on Earth. The strength of Martian gravity being about 38% of that on Earth, a kicked-up grain is accelerated by the wind roughly seven times longer than an equivalent grain on Earth. When feedback effects are included, Martian sand grains attain ten times the speed of their terrestrial counterparts. The researchers have found that certain observed dune fields are consistent with wind friction

speeds of about 3 meters per second. Other fields, they've found, can be explained (in their model) only under the assumption of a bimodal wind regime, with the wind alternating between two directions (see Figure 4).

The study of dunes, Herrmann points out, involves all three components of modern science: theory, computation, and experiment. Field measurements, he says, are vital. Given where dunes tend to be found, experiments are also hot, dirty, grueling work. Herrmann recalls an urgent call from Parteli's mother, shortly after Parteli returned from a two-week trek through South America, begging Herrmann not to send her son on the next field trip. Parteli, it seems, had mentioned that for his next assignment he would be studying the dunes on Mars.

Barry A. Cipra is a mathematician and writer based in Northfield, Minnesota.




Figure 3. Simulations show various possible outcomes when a smaller barchan overtakes a larger one, depending on their relative sizes: coalescence (a), "breeding" (b and c), "budding" (d), and solitary wave behavior (e). The height of the larger barchan is 5 meters; the heights of the smaller barchans are 1.8, 1.9, 2.2, 2.6, and 3.1 meters, respectively. Time is given in months. From "Breeding and Solitary Wave Behavior of Dunes," by Oscar Durán, Veit Schwämmle, and Hans Herrmann.

GAMS

OPTIMIZATION

High Level Modeling

The General Algebraic Modeling System (GAMS) is a **high level modeling system** for mathematical programming problems. GAMS is tailored for both developing research-driven rapid prototype models all the way to large-scale commercial applications on a wide variety of platforms.

Multiple Model Types

GAMS allows the formulation of models in **many problem classes**, including linear (LP) and mixed integer linear (MIP), nonlinear (NLP) and mixed integer nonlinear (MINLP), mixed complementarity (MCP), stochastic LP, as well as quadratic programming (QCP).

State-of-the-Art Solvers

GAMS incorporates all major commercial and academic **state-of-the-art solution technologies** for a broad range of problem types, including a range of **global solvers**.



Download a free demo at www.gams.com

Email: sales@gams.com
Phone: (202) 342-0180



IMA INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS

University of Minnesota, Minneapolis, Minnesota

Associate Director

The Institute for Mathematics and its Applications (IMA) seeks a person to serve as Associate Director for two years beginning in August 2008.

The IMA is located on the Minneapolis campus of the University of Minnesota. Since its founding in 1982 as the result of a national competition, the IMA has been a major national institute with the mission of fostering interdisciplinary research which links high caliber mathematics with important problems from other disciplines, technology, industry, and society, and of strengthening the talent base engaged in mathematical research applied to or relevant to such problems. The IMA runs visitor programs involving around 1,200 scientists per year, with an annual budget of about \$6M. The major source of funding of the IMA is the National Science Foundation which has funded the IMA through 2010. The IMA's many programs and activities are documented at its web site, www.ima.umn.edu.

The Associate Director will work with the Director and Deputy Director, the IMA staff, and program organizers on a variety of IMA activities, especially program implementation, participant applications, publicity and special projects. A successful candidate should have a PhD in mathematics or a related field and a successful record in and broad view of mathematical research, a commitment to the IMA mission, and administrative or organizational experience. The Associate Director will work in the exceptional scientific environment provided by the IMA, and will contribute to shaping the international research agenda. Research activity is compatible with the position, and a visiting faculty position at the University of Minnesota may be arranged if appropriate.

Applicants should send a CV, a statement of their interest in the position and other relevant information not contained in the CV, and have at least three letters of reference sent. All materials should be sent by email to search@ima.umn.edu and inquiries can be directed to the same address. Review of applications will begin on January 14, 2008. Applications will be accepted until the position is filled.

The University of Minnesota is an equal opportunity educator and employer.



The IMA is an NSF funded Institute

